

Poincaré inequalities and rigidity for actions on Banach spaces

Piotr W. Nowak

Abstract

The aim of this paper is to extend the framework of the spectral method for proving property (T) to the class of reflexive Banach spaces and present a condition implying that every affine isometric action of a given group G on a reflexive Banach space X has a fixed point. This last property is a strong version of Kazhdan's property (T) and is equivalent to the fact that $H^1(G, \pi) = 0$ for every isometric representation π of G on X . The condition is expressed in terms of p -Poincaré constants and we provide examples of groups, which satisfy such conditions and for which $H^1(G, \pi)$ vanishes for every isometric representation π on an L_p space for some $p > 2$. Our methods allow to estimate such a p explicitly and yield several interesting applications. In particular, we obtain quantitative estimates for vanishing of 1-cohomology with coefficients in uniformly bounded representations on a Hilbert space. We also give lower bounds on the conformal dimension of the boundary of a hyperbolic group in the Gromov density model.

Keywords. Poincaré inequality; Kazhdan's property (T); affine isometric action; 1-cohomology.

1 Introduction

Kazhdan's property (T) is a powerful rigidity property of groups with numerous applications and several characterizations. In this article we focus on the following description of property (T): *a group G has property (T) if and only if every affine isometric action of G on the Hilbert space has a fixed point.* This characterization can be rephrased as the cohomological condition $H^1(G, \pi) = 0$, for every unitary representation π of G . A generalization of property (T) to other Banach spaces is then straightforward: we are interested in conditions implying that every affine isometric action of a given group on a given Banach space has a fixed point. Such rigidity properties for actions on Banach spaces, as

Institute of Mathematics of the Polish Academy of Sciences, Śniadeckich 8, 00-656 Warszawa, Poland
Institute of Mathematics, University of Warsaw, Banacha 2, 02-097 Warszawa, Poland. e-mail: pnowak@mimuw.edu.pl

Mathematics Subject Classification (2010): Primary 22D12; Secondary 46B03

well as other generalizations of property (T), and their applications, were studied earlier in [2, 12, 8, 21].

One very successful method of proving property (T) is through spectral conditions on links of vertices of complexes acted upon by a group. Variations of such conditions were studied in [3, 9, 10, 14, 17, 19, 33, 38, 39, 35, 36] in the context of Hilbert spaces and non-positively curved spaces. Given a group G acting on a 2-dimensional simplicial complex, one considers the link of a vertex. This link is a finite graph. If for every vertex, the first positive eigenvalue of the discrete Laplacian is strictly larger than $1/2$, then G has property (T).

The main purpose of this work is to extend the framework of the spectral method, and some of the rigidity results, beyond Hilbert spaces. Our main result provides such a framework for the class of reflexive Banach spaces. The difficulty lies in the fact, that in the Hilbert space case the spectral method relies heavily on orthogonality, in particular self-duality of Hilbert spaces. When passing to other Banach spaces, dual spaces of certain Banach spaces and of their subspaces have to be identified, and this is often a difficult task. We show, that when the representation is isometric such computations are possible and we can use duality effectively.

We focus on link graphs constructed using generating sets of a group, as in [39]. For a finite, symmetric generating set S not containing the identity element the vertices of the link graph $\mathcal{L}(S)$ are the elements of S ; generators s and t are connected by an edge if $s^{-1}t$ is a generator. We will also assume that the graph is equipped with a weight ω on the edges.

Let X be a Banach space and denote by $\kappa_p(S, X)$ be the optimal constant in the p -Poincaré inequality for the link graph $\mathcal{L}(S)$ of G and the norm of X ,

$$\sum_{s \in S} \|f(s) - Af\|_X^p \deg_\omega(s) \leq \kappa_p^p \sum_{s \sim t} \|f(s) - f(t)\|_X^p \omega(s, t),$$

where Af is the mean value of f . When $X = L_2$, the constant $\kappa_2(S, L_2) = \kappa_2(S, \mathbb{R})$ can be expressed in terms of the first eigenvalue of the discrete Laplacian.

Our main result shows that sufficiently small constants in Poincaré inequalities for the graph $\mathcal{L}(S)$ imply the required cohomological vanishing. Given a number $1 < p < \infty$ we denote by p^* its adjoint index, satisfying $\frac{1}{p} + \frac{1}{p^*} = 1$.

Theorem 1.1. *Let X be a reflexive Banach space and let G be a group generated by a finite, symmetric set S , not containing the identity element. If the link graph $\mathcal{L}(S)$ is connected and for some $1 < p < \infty$ the associated Poincaré constants satisfy*

$$\max \left\{ 2^{-\frac{1}{p}} \kappa_p(S, X), 2^{-\frac{1}{p^*}} \kappa_{p^*}(S, X^*) \right\} < 1,$$

then

$$H^1(G, \pi) = 0,$$

for every isometric representation π of G on X .

Clearly, by reflexivity, the same conclusion holds for actions on X^* . Interestingly, the roles of the two constants in the proof of the above theorem are not symmetric.

We apply Theorem 1.1 to L_p spaces. The interesting case is $p > 2$. Indeed, when $1 < p \leq 2$, affine isometric actions exhibit the same behavior as for the Hilbert space: G has property (T) if and only if any affine action on an L_p space for $1 < p \leq 2$ has a fixed point [2]. Also, G admits a metrically proper affine isometric action on the Hilbert space (i.e., is a-T-menable) if and only if it admits such an action on any $L_p[0, 1]$ for $1 < p \leq 2$ [28] (see corrected version [29]). This last property is a strong negation of the existence of a fixed point.

Fixed point properties for groups acting on L_p spaces for $p > 2$ are difficult to prove and only a handful of results are known:

1. higher rank algebraic groups and their lattices have fixed points for every affine isometric action on L_p -spaces for all $p > 1$ [2];
2. in [23] it was proved that $\mathrm{SL}_n(\mathbb{Z}[x_1, \dots, x_k])$ has fixed points for every affine isometric on L_p for every $p > 1$ and $n \geq 4$;
3. Naor and Silberman [24] showed that Gromov's random groups, containing (in a certain weak sense) expanders in their Cayley graphs, have a fixed point for affine isometric actions on any L_p for $p > 1$;
4. a general argument due to Fisher and Margulis (see the proof in [2]) shows, that for every property (T) group G there exists a constant $\varepsilon = \varepsilon(G) > 0$ such that any affine isometric action on L_p for $p \in [2, 2 + \varepsilon)$, has a fixed point. However, their argument does not give any control over ε .

On the other hand there are also groups which have property (T) but act without fixed points on L_p spaces. One example is furnished by $\mathrm{Sp}(n, 1)$, which has property (T) but has non-vanishing L_p -cohomology for $p > 4n + 2$, by a result of Pansu [32]. It is also known that there exist hyperbolic groups which have property (T). Nevertheless, Bourdon and Pajot [5] showed that for every hyperbolic group G and sufficiently large $p > 2$ there is an affine isometric action on $\ell_p(G)$, whose linear part is the regular representation and which does not have a fixed point. Moreover, Yu [37] showed that every hyperbolic group admits a proper, affine isometric action on $\ell_p(G \times G)$ for all sufficiently large $p > 2$ (see also [27] for another construction). We refer to [30] for a recent survey.

The techniques we use to establish the fixed point properties are different from the ones used previously for general Banach spaces. In particular, we do not need the Howe-Moore property to prove our results. This representation theoretic property was necessary in [2, 23]. The expected outcome is also slightly different, as our methods are not expected to give fixed points on L_p for all $p > 1$. One reason is that the p -Poincaré constants usually increase above $2^{1/p}$ as p grows to infinity. The second reason is that the main result applies to random hyperbolic groups, which, as remarked earlier, act without fixed points on L_p -spaces for $p > 2$ sufficiently large. Using our approach we obtain the appropriate vanishing of cohomology $H^1(G, \pi)$ for isometric representations π on L_p -spaces with $p \in [2, 2 + c)$, where the value of c depends on the group and can be estimated explicitly. Finally, we point out that our techniques and the Poincaré inequalities we use, are all linear, in contrast to the non-linear approach used e.g., in [17, 35]. Linearity allows to

use interpolation methods effectively and also to obtain additional information about the structure of cohomology in the presence of spectral gaps.

To apply Theorem 1.1 we need to estimate p -Poincaré constants for $p > 2$. Even in classical settings, such as convex domains in \mathbb{R}^n , estimates exist but exact values of p -Poincaré constants are not known, except a few special cases. The situation is even worse for finite graphs, where very few estimates are known for cases other than $p = 1, 2$. Here we consider the family of A_2 -groups, indexed by powers of primes. These groups were introduced and studied in [6, 7]. For every q , the group G_q has a generating sets whose link graph is the incidence graph of the finite projective plane over the field \mathbb{F}_q . Spectra of such graphs were computed in [11] and give, in particular, the exact value of the Poincaré constant $\kappa_2(S, \mathbb{R})$. We use this fact to estimate $\kappa_p(S, L_p)$ for these graphs, which allows to obtain for each q a number c_q such that any affine isometric action of G on any L_p has a fixed point for $p \in [2, 2 + c_q)$. The explicit estimates of c_q are given in Theorem 5.1.

As mentioned earlier, our results apply to random hyperbolic groups, more precisely, to random groups in the Gromov density model with densities $1/3 < d < 1/2$, and yield important consequences. These groups are hyperbolic and have Kazhdan's property (T) with overwhelming probability [39, 20]. Using our methods we give lower bounds on p for which fixed points exists for all isometric actions on any L_p -space. A connection with the conformal dimension arises through the work of Bourdon and Pajot [5] and allows us to give a lower bound on the conformal dimension of a boundary of a random hyperbolic group, using an associated link graph, see Section 6. The problem of estimating the conformal dimension of random hyperbolic groups was posed by Gromov [16, 9.B (g)] and Pansu [31, IV.b].

Our methods also apply to affine actions, whose linear part is a uniformly bounded representation on a Hilbert space. More precisely, we show that $H^1(G, \pi) = 0$, whenever π is a uniformly bounded representation with norms of all operators bounded by a constant, which depends on the group but is close to $\sqrt{2}$ in many cases, see Theorems 5.5 and 6.3. The question of extending property (T) in the form of cohomological vanishing from unitary to uniformly bounded representation is a well-known open question. In particular, Shalom conjectures that for every hyperbolic group there exists a uniformly bounded representation with a proper cocycle. The case of $\mathrm{Sp}(n, 1)$ is an unpublished result of Shalom.

Finally, we present other applications. We improve the differentiability class of diffeomorphic actions on the circle in the rigidity theorem in [25, 26] and estimate eigenvalues of the discrete p -Laplacian on finite quotients of groups using Kazhdan-type constants.

Contents

| | | |
|----------|--|----------|
| 1 | Introduction | 1 |
| 2 | Actions on Banach spaces | 5 |
| 2.1 | Generating sets and link graphs | 5 |
| 2.2 | Isometric representations and associated Banach spaces | 6 |
| 2.3 | Duality for $C_-^{(1,p)}(G, \pi)$ | 7 |

| | |
|--|-----------|
| <i>JEMS template</i> | 5 |
| 2.4 The operator δ | 9 |
| 2.5 The operators D , L , and d | 9 |
| 2.6 Sufficient conditions for vanishing of cohomology | 11 |
| 3 Poincaré inequalities associated to norms | 12 |
| 3.0.1 Hilbert spaces | 12 |
| 3.0.2 L_p -spaces, $1 \leq p < \infty$ | 12 |
| 3.0.3 Direct sums | 13 |
| 3.0.4 The case $p = \infty$ | 13 |
| 3.0.5 Behavior under isomorphisms | 13 |
| 4 Vanishing of cohomology | 13 |
| 4.1 An inequality for κ_p and δ_π^* | 13 |
| 4.2 Proof of the main theorem | 14 |
| 5 \tilde{A}_2 groups | 16 |
| 6 Hyperbolic groups | 18 |
| 7 Other Applications | 20 |
| 7.1 Actions on the circle | 20 |
| 7.2 The finite-dimensional case and the p -Laplacian | 21 |

2 Actions on Banach spaces

2.1 Generating sets and link graphs

Let G denote a discrete group generated by a finite symmetric set $S = S^{-1}$. Let $\mathcal{L}(S)$ denote the following graph, called the link graph of S . The vertices are given by $\mathcal{V} = S$. Two vertices $s, t \in S$ are connected by an edge, denoted $s \sim t$, if and only if $s^{-1}t \in S$ and $t^{-1}s \in S$.

The set E is defined as follows:

$$E = \{(s, t) \in S \times S : s^{-1}t \in S\}.$$

Note that E can be viewed as the set of oriented edges and in E every edge is counted twice.

A *weight* on $\mathcal{L}(S)$ is a function $\omega : E \rightarrow (0, \infty)$, such that $\omega(s, t) = \omega(t, s)$, for every $s, t \in S$. Given a weight on the link graph, the associated *degree* of a vertex $r \in S$ is defined to be

$$\deg_\omega(s) = \sum_{t, t \sim s} \omega(t, s).$$

A weight ω on a link graph $\mathcal{L}(S)$ is *admissible* if it satisfies

1. $\deg_\omega(s) = \deg_\omega(s^{-1})$, and

$$2. \deg_\omega(r) = \sum_{(s,t): s^{-1}t=r} \omega(s, t),$$

for every $r, s, t \in S$. Note that

$$\sum_{s \in S} \deg_\omega(s) = \omega(E).$$

Throughout the article we consider only admissible weights on link graphs of generating sets.

2.2 Isometric representations and associated Banach spaces

Let X be a Banach space equipped with a norm $\|\cdot\|_X$. We assume throughout that X is reflexive and that $\pi : G \rightarrow B(X)$ is a representation of G into the bounded invertible operators on X . Let X^* denote the continuous dual of X , with its standard norm. X^* is naturally equipped with the adjoint representation of G , $\bar{\pi} : G \rightarrow B(X^*)$,

$$\bar{\pi}_g = \pi_{g^{-1}}^*.$$

Throughout we fix $1 < p < \infty$. The value of p will be chosen later depending on the context. We denote by p^* the adjoint index, satisfying $\frac{1}{p} + \frac{1}{p^*} = 1$, and by L_p the space $L_p(\mu)$ for any measure μ (our results apply with no assumptions on the measure). We also use \simeq to denote an isomorphism and \cong to denote an isometric isomorphism of Banach spaces.

Define the Banach space $C^{(0,p)}(G, \pi)$ to be the linear space X , with the norm

$$\|v\|_{(0,p)} = \omega(E)^{\frac{1}{p}} \|v\|_X.$$

Let $\langle \cdot, \cdot \rangle_X$ denote the natural pairing between X and X^* . The pairing between $C^{(0,p)}(G, \pi)$ and $C^{(0,p^*)}(G, \bar{\pi})$ is given by

$$\langle v, w \rangle_0 = \omega(E) \langle v, w \rangle_X.$$

Then $C^{(0,p^*)}(G, \bar{\pi})$ is the dual space of $C^{(0,p)}(G, \pi)$.

We define $C^{(1,p)}(G, \pi)$ to be the finite direct sum $\bigoplus_{s \in S} X$, with the norm given by

$$\|f\|_{(1,p)} = \left(\sum_{s \in S} \|f(s)\|_X^p \deg_\omega(s) \right)^{\frac{1}{p}}.$$

The dual of $C^{(1,p)}(G, \pi)$ is $C^{(1,p^*)}(G, \bar{\pi})$, via the pairing

$$\langle f, \phi \rangle_1 = \sum_{s \in S} \langle f(s), \phi(s) \rangle_X \deg_\omega(s),$$

for $f \in C^{(1,p)}(G, \pi)$ and $\phi \in C^{(1,p^*)}(G, \bar{\pi})^*$.

Define an operator Q_π on $C^{(1,p)}(G, \pi)$,

$$Q_\pi f(s) = \pi_s f(s^{-1}).$$

A similar operator $Q_{\bar{\pi}}$ is defined on $C^{(1,p^*)}(G, \bar{\pi})$. The following is straightforward to verify.

Lemma 2.1. *The operator Q_π is an involution, satisfying $Q_\pi^* = Q_{\bar{\pi}}$.*

Consider the following subspaces of $C^{(1,p)}(G, \pi)$, defined as eigenspaces of Q_π :

$$C_+^{(1,p)}(G, \pi) = \{f \in C^{(1,p)}(G, \pi) : f = Q_\pi f\},$$

and

$$C_-^{(1,p)}(G, \pi) = \{f \in C^{(1,p)}(G, \pi) : f = -Q_\pi f\}.$$

Lemma 2.2. *For any $1 < p < \infty$ we have $C^{(1,p)}(G, \pi) = C_+^{(1,p)}(G, \pi) \oplus C_-^{(1,p)}(G, \pi)$.*

Proof. We define two bounded operators: $P_\pi^+ : C^{(1,p)}(G, \pi) \rightarrow C_+^{(1,p)}(G, \pi)$,

$$P_\pi^+ = \frac{I + Q_\pi}{2},$$

and $P_\pi^- : C^{(1,p)}(G, \pi) \rightarrow C_-^{(1,p)}(G, \pi)$,

$$P_\pi^- = \frac{I - Q_\pi}{2}.$$

Clearly $P_\pi^+ + P_\pi^- = I$. Additionally, $C_+^{(1,p)}(G, \pi) = \ker P_\pi^- = \text{im } P_\pi^+$ and $C_-^{(1,p)}(G, \pi) = \ker P_\pi^+ = \text{im } P_\pi^-$. Indeed, we have

$$\pi_{s^{-1}}(P_\pi^+ f(s)) = \frac{\pi_{s^{-1}} f(s) + f(s^{-1})}{2} = P_\pi^+ f(s^{-1}).$$

Finally, P_π^+ restricted to $C_+^{(1,p)}(G, \pi)$ and P_π^- restricted to $C_-^{(1,p)}(G, \pi)$ are identity operators, so that P_π^+ and P_π^- are projections onto the required subspaces. \square

We now analyze the structure of $C^{(1,p)}(G, \pi)$ in relation to the one of $C^{(1,p^*)}(G, \bar{\pi})$.

2.3 Duality for $C_-^{(1,p)}(G, \pi)$

The dual of $C^{(1,p)}(G, \pi)$ is $C^{(1,p^*)}(G, \bar{\pi})$. Let $P_{\bar{\pi}}^+ : C^{(1,p^*)}(G, \bar{\pi}) \rightarrow C_+^{(1,p^*)}(G, \bar{\pi})$ and $P_{\bar{\pi}}^- : C^{(1,p^*)}(G, \bar{\pi}) \rightarrow C_-^{(1,p^*)}(G, \bar{\pi})$ denote similar projections as above on the dual level.

Lemma 2.3. *We have $P_{\bar{\pi}}^+ = (P_\pi^+)^*$ and $P_{\bar{\pi}}^- = (P_\pi^-)^*$.*

Proof. Let $f \in C^{(1,p)}(G, \pi)$ and $\phi \in C^{(1,p^*)}(G, \bar{\pi})$. Then

$$(P_\pi^-)^* = \frac{1}{2}(I - Q_\pi)^* = \frac{1}{2}(I - Q_{\bar{\pi}}) = P_{\bar{\pi}}^-.$$

Similarly for P_π^+ . \square

Lemma 2.4. *We have the following isomorphisms: $C_-^{(1,p)}(G, \pi)^* \simeq C_-^{(1,p^*)}(G, \bar{\pi})$ and $C_+^{(1,p)}(G, \pi)^* \simeq C_+^{(1,p^*)}(G, \bar{\pi})$.*

Proof. Consider $f \in C_-^{(1,p)}(G, \pi)$ and let $\phi \in C^{(1,p^*)}(G, \bar{\pi})$. Then

$$\langle f, \phi \rangle_1 = \langle -Q_\pi f, \phi \rangle_1 = \langle f, -Q_\pi \phi \rangle_1.$$

Therefore,

$$2\langle f, P_\pi^+ \phi \rangle_1 = 0,$$

which shows that $C_+^{(1,p^*)}(G, \bar{\pi})$ annihilates $C_-^{(1,p)}(G, \pi)$.

Conversely, if $\phi \in C^{(1,p^*)}(G, \bar{\pi})$ annihilates $C_-^{(1,p)}(G, \pi)$, then

$$\langle P_\pi^- f, \phi \rangle_1 = \langle f, P_\pi^- \phi \rangle_1 = 0$$

for every $f \in C^{(1,p)}(G, \pi)$. Consequently, $P_\pi^- \phi = 0$ and $\phi = P_\pi^+ \phi$, which means it belongs to $C_+^{(1,p^*)}(G, \bar{\pi})$. Thus,

$$C_-^{(1,p)}(G, \pi)^* \cong C^{(1,p)}(G, \bar{\pi}) / C_+^{(1,p^*)}(G, \bar{\pi}) \simeq C_-^{(1,p^*)}(G, \bar{\pi}).$$

Other cases are proved similarly. \square

However, in order to identify the dual of $C_-^{(1,p)}(G, \pi)$ an isomorphism is not sufficient, we need an isometric isomorphism instead. For a representation π , $C_-^{(1,p)}(G, \pi)^*$ is in general not isometrically isomorphic to $C_-^{(1,p^*)}(G, \bar{\pi})$. However, it turns out that this additional property holds when the representation π is isometric.

Theorem 2.5. *Assume that π_s is an isometry for every $s \in S$. Then we have the following isometric isomorphisms: $C_-^{(1,p)}(G, \pi)^* \cong C_-^{(1,p^*)}(G, \bar{\pi})$ and $C_+^{(1,p)}(G, \pi)^* \cong C_+^{(1,p^*)}(G, \bar{\pi})$.*

Proof. Consider $C^{(1,p^*)}(G, \bar{\pi}) / C_+^{(1,p^*)}(G, \bar{\pi})$, which consists of cosets $[\phi] = C_+^{(1,p^*)}(G, \bar{\pi}) + \phi$, for $\phi \in C^{(1,p^*)}(G, \bar{\pi})$. We need to show that for each such coset N , $\inf \{ \|\phi\| : N = [\phi] \}$ is attained when $\phi \in C_-^{(1,p^*)}(G, \bar{\pi})$.

For $\phi \in C_-^{(1,p^*)}(G, \bar{\pi})$ and $\psi \in C_+^{(1,p^*)}(G, \bar{\pi})$, we have

$$\|\phi + \psi\|_{(1,p^*)} = \| -Q_\pi \phi + Q_\pi \psi \|_{(1,p^*)} = \|\phi - \psi\|_{(1,p^*)},$$

since the involution Q_π is an isometry, whenever π , or equivalently $\bar{\pi}$, is an isometric representation. Now consider the coset $[\phi]$ for $\phi \in C_-^{(1,p^*)}(G, \bar{\pi})$ and consider another element, $\zeta \in C^{(1,p^*)}(G, \bar{\pi})$, such that $\zeta - \phi \in C_+^{(1,p^*)}(G, \bar{\pi})$, so that $\zeta = \phi + \psi$, for some $\psi \in C_+^{(1,p^*)}(G, \bar{\pi})$. This implies

$$\|\phi\|_{(1,p^*)} \leq \frac{\|\phi - \psi\|_{(1,p^*)} + \|\phi + \psi\|_{(1,p^*)}}{2} = \|\zeta\|_{(1,p^*)},$$

which proves the claim. \square

This last statement allows us to identify $C_-^{(1,p)}(G, \pi)^*$ with $C_-^{(1,p^*)}(G, \bar{\pi})$ for isometric representations and is crucial in the proof of the main theorem.

2.4 The operator δ

We define the operator $\delta_\pi : C^{(0,p)}(G, \pi) \rightarrow C_-^{(1,p)}(G, \pi)$ by the formula

$$\delta_\pi v(s) = v - \pi_s v.$$

Theorem 2.5 allows to express the adjoint of δ_π in a way, which is convenient for calculations. We have the following explicit formula for δ_π^* .

Lemma 2.6. *The operator $\delta_\pi^* : C_-^{(1,p^*)}(G, \bar{\pi}) \rightarrow C^{(0,p^*)}(G, \bar{\pi})$ is given by*

$$\delta_\pi^* \phi = 2 \sum_{s \in S} \phi(s) \frac{\deg_\omega(s)}{\omega(E)}. \quad (2.1)$$

Proof.

$$\begin{aligned} \langle \delta_\pi v, \phi \rangle_1 &= \sum_{s \in S} \langle v - \pi_s v, \phi(s) \rangle_X \deg_\omega(s) \\ &= \sum_{s \in S} (\langle v, \phi(s) \rangle_X - \langle v, \bar{\pi}_{s^{-1}} \phi(s) \rangle_X) \deg_\omega(s) \\ &= \sum_{s \in S} (\langle v, \phi(s) \rangle_X + \langle v, \phi(s^{-1}) \rangle_X) \deg_\omega(s) \\ &= \left\langle v, 2 \sum_{s \in S} \phi(s) \frac{\deg_\omega(s)}{\omega(E)} \right\rangle_0. \end{aligned}$$

□

It is now clear that δ_π^* admits a continuous extension to the space $C^{(1,p^*)}(G, \bar{\pi})$, defined by the right hand side of the formula (2.1).

2.5 The operators D , L , and d

We define the Banach space,

$$C^{(2,p)}(G, \pi) = \left\{ \eta \in \bigoplus_{(s,t) \in E} X : \eta(s,t) = -\eta(t,s) \right\},$$

equipped with the norm

$$\|\eta\|_{(2,p)} = \left(\sum_{(s,t) \in E} \|\eta(s,t)\|_X^p \omega(s,t) \right)^{\frac{1}{p}}.$$

We also define operators $D, L_\pi : C_-^{(1,p)}(G, \pi) \rightarrow C^{(2,p)}(G, \pi)$ by the formulas

$$Df(s,t) = f(t) - f(s),$$

$$L_\pi f(s, t) = \pi_s f(s^{-1}t).$$

Then the operator d_π is defined by

$$d_\pi = L_\pi - D.$$

Similarly, define \overline{D} , $L_{\overline{\pi}}$ and $d_{\overline{\pi}}$ for the adjoint representation.

Lemma 2.7. *Let π be an isometric representation. The operator L_π is an isometry onto its image. Consequently, D is an isometry as well when restricted to $\ker d_\pi$. (The same claim holds for $L_{\overline{\pi}}$ and \overline{D} , restricted to $\ker d_{\overline{\pi}}$).*

Proof. By direct calculation,

$$\begin{aligned} \|L_\pi f\|_{(2,p)}^p &= \sum_{(s,t) \in E} \|\pi_s f(s^{-1}t)\|_X^p \omega(s, t) \\ &= \sum_{s \in S} \|f(s)\|_X^p \deg_\omega(s) \\ &= \|f\|_{(1,p)}^p. \end{aligned}$$

□

The kernel of the operator \overline{D} consists of the constant functions on S , which is a complemented subspace of $C_-^{(1,p)}(G, \pi)$. The projection onto this subspace is given by

$$\overline{M}\phi(s) = \sum_{s \in S} \phi(s) \frac{\deg_\omega(s)}{\omega(E)}.$$

Note that for $\phi \in C_-^{(1,p^*)}(G, \overline{\pi})$ we have

$$\overline{M}\phi(s) = \frac{1}{2} \delta_\pi^* \phi,$$

for every $s \in S$.

Lemma 2.8. *Let $\phi \in C_-^{(1,p^*)}(G, \overline{\pi})$. Then $\|\overline{M}\phi\|_{(1,p^*)} = \frac{1}{2} \|\delta_\pi^* \phi\|_{(0,p^*)}$.*

Proof. We have the following equalities:

$$\begin{aligned} \|\overline{M}\phi\|_{(1,p^*)}^{p^*} &= \sum_{s \in S} \left\| \frac{\delta_\pi^* \phi}{2} \right\|_X^{p^*} \deg_\omega(s) \\ &= \frac{1}{2^{p^*}} \|\delta_\pi^* \phi\|_X^{p^*} \left(\sum_{s \in S} \deg_\omega(s) \right) \\ &= \frac{1}{2^{p^*}} \|\delta_\pi^* \phi\|_0^{p^*}. \end{aligned}$$

□

2.6 Sufficient conditions for vanishing of cohomology

Given a group G , the 1-cocycles associated to π are functions $b : G \rightarrow X$ satisfying the cocycle condition,

$$b_g = \pi_g b_h + b_g,$$

for every $g, h \in G$. The coboundaries are those cocycles which are of the form

$$b_g = v - \pi_g v$$

for some $v \in X$ and all $g \in G$. The first cohomology of G with coefficients in π is defined to be $H^1(G, \pi) = \text{cocycles} / \text{coboundaries}$.

An affine action of G on X is defined as

$$A_g v = \pi_g v + b_g,$$

where π is called the linear part of the action and b is a cocycle. Vanishing of cohomology $H^1(G, \pi)$ is equivalent to the existence of a fixed point for any affine action with linear part π . We refer to [4] for background on cohomology and affine actions.

The reader can easily verify the following lemma.

Lemma 2.9. $\text{image}(\delta_\pi) \subseteq \ker d_\pi$.

This fact allows to formulate the following sufficient condition for the fixed point property for affine actions on X .

Proposition 2.10. *If the image of δ_π is equal to $\ker d_\pi$, then $H^1(G, \pi) = 0$.*

Proof. Let $b : G \rightarrow X$ be a 1-cocycle for π and let b' denote the restriction of b to the generating set S . The cocycle condition implies that $b' \in C_-^{(1,p)}(G, \pi)$ and, furthermore, that $b' \in \ker d_\pi$. If δ_π is onto $\ker d_\pi$, then $b' = \delta_\pi v$ for some $v \in X$. Since b is trivial on the generators, we conclude that b is trivial. \square

It is important to remark that the technical details here are slightly different than in [39], where the original condition in terms of almost invariant vectors is deduced, and one needs to use the Delorme-Guichardet theorem to obtain cohomological vanishing. The above argument allows to bypass the use of the Delorme-Guichardet theorem and obtain vanishing of cohomology directly.

Note that the image of δ_π is always properly contained in $C_-^{(1,p)}(G, \pi)$. By the open mapping theorem we also have the following

Corollary 2.11. *Assume π does not have invariant vectors. If δ is onto $\ker d$ then there is a constant $K > 0$ such that*

$$\sup_{s \in S} \|v - \pi_s v\|_X \geq K \|v\|_X,$$

for every $v \in X$.

The constant K in the above statement can be viewed as a version of Kazhdan constant for isometric representations of G on X .

3 Poincaré inequalities associated to norms

Consider a weighted, finite graph $\Gamma = (\mathcal{V}, \mathcal{E})$, a number $p \geq 1$ and a Banach space X . The p -Poincaré inequality for Γ and the norm of X is the inequality

$$\left(\sum_{x \in \mathcal{V}} \|f(x) - Af\|_X^p \deg_\omega(x) \right)^{\frac{1}{p}} \leq \kappa_p \left(\sum_{x \sim y} \|f(x) - f(y)\|_X^p \omega(x, y) \right)^{\frac{1}{p}}, \quad (3.1)$$

for all functions $f : \mathcal{V} \rightarrow X$, where $Af = \frac{1}{2\omega(\mathcal{E})} \sum_{x \in \mathcal{V}} f(x) \deg_\omega(x)$. On a finite graph, the inequality (3.1) is always satisfied for some $\kappa_p > 0$.

Definition 3.1. Let $\mathcal{L}(S)$ be a link graph of a generating set S , with weight ω . For a Banach space X and a number $1 < p < \infty$ we define the constant $\kappa_p(S, X)$ of $\mathcal{L}(S)$ by setting

$$\kappa_p(S, X) = \inf \kappa_p,$$

where the infimum is taken over all κ_p , for which inequality (3.1) holds.

We will omit the reference to ω in the notation for κ .

3.0.1 Hilbert spaces

When $X = L_2$ is the Hilbert space this constant is related to the smallest positive eigenvalue λ_1 of the Laplacian on the graph as follows:

$$\kappa_2(S, L_2) = \sqrt{\lambda_1^{-1}},$$

since the latter can be defined via the variational expression and the Rayleigh quotient.

3.0.2 L_p -spaces, $1 \leq p < \infty$

Let (Y, μ) be any measure space and for $X = \mathbb{R}$ consider a p -Poincaré inequality

$$\sum_{x \in \mathcal{V}} |f(x) - Af|^p \deg_\omega(x) \leq \kappa_p^p \sum_{x \sim y} |f(x) - f(y)|^p \omega(x, y) \quad (3.2)$$

on a finite graph. By integrating over Y with respect to μ we obtain

$$\sum_{x \in \mathcal{V}} \|f(x) - Af\|_{L_p}^p \deg_\omega(x) \leq \kappa_p^p \sum_{x \sim y} \|f(x) - f(y)\|_{L_p}^p \omega(x, y),$$

for any $f : \mathcal{V} \rightarrow L_p$. This gives

$$\|f - Af\|_{(1,p)} \leq \kappa_p \|\nabla f\|_{(2,p)},$$

so that $\kappa_p(S, L_p)$ is equal to $\kappa_p(S, \mathbb{R})$ in the inequality (3.2).

3.0.3 Direct sums

More generally, consider an ℓ_p -direct sum $X = \left(\bigoplus_{s \in S} X_s\right)_p$ of Banach spaces $\{X_s\}_{s \in S}$. A similar argument as above shows that $\kappa_p(S, X) \leq \sup_{s \in S} \kappa_p(S, X_s)$.

3.0.4 The case $p = \infty$

Consider $s, s^{-1} \in S$ and choose $x \in X$ such that $\|x\|_X = 1$. Let d_S denote the path metric on $\mathcal{L}(S)$. Define $f : \Gamma \rightarrow \mathbb{R}$ by the formula

$$f(t) = \begin{cases} \left(1 - 2 \frac{d_S(s, t)}{d_S(s, s^{-1})}\right) x & \text{if } d_S(s, t) \leq \frac{d_S(s, s^{-1})}{2}, \\ \left(-1 + 2 \frac{d_S(s, t)}{d_S(s, s^{-1})}\right) x & \text{if } d_S(s^{-1}, t) \leq \frac{d_S(s, s^{-1})}{2}, \\ 0 & \text{if } d_S(s, t) > \frac{d_S(s, s^{-1})}{2} \text{ and } d_S(s^{-1}, t) > \frac{d_S(s, s^{-1})}{2}. \end{cases}$$

For such f we have $\|f\|_{(X, \infty)} = 1$ and $Af = 0$, however $\|Df\|_{(X, \infty)} = \frac{1}{d_S(s, s^{-1})}$. Thus we have

$$\kappa_\infty(G, \pi) \geq \max_{s \in S} d_S(s, s^{-1})$$

and for sufficiently large S , the above Poincaré constant is at least 1. Additionally, for any $\varepsilon > 0$ there exists a sufficiently large $p < \infty$, such that the norms $\|f\|_{(1, p)}$ and $\|f\|_{(1, \infty)}$ are ε -close. For a sufficiently small $\varepsilon > 0$ and the corresponding p as above, we also have $2^{-\frac{1}{p}} \kappa_p(S, X) \geq 1$.

3.0.5 Behavior under isomorphisms

Let $T : X \rightarrow Y$ be an isomorphism of Banach spaces X, Y , satisfying $\|x\|_X \leq \|Tx\|_Y \leq L\|x\|_X$ for every $x \in X$. Then $\kappa_p(G, \pi) \leq L\kappa_p(G, Y)$.

4 Vanishing of cohomology

4.1 An inequality for κ_p and δ_π^*

Note that since in E each edge of $\mathcal{L}(S)$ is counted twice, we have $\|Df\|_{(2, p)} = 2^{\frac{1}{p}} \|\nabla f\|_{\ell_p(S, X)}$ and $Mf = Af$. The following result describes the relation between Poincaré constants and the operator δ_π^* .

Theorem 4.1. *The inequality*

$$2 \left(1 - 2^{-\frac{1}{p^*}} \kappa_{p^*}(S, X)\right) \|\phi\|_{(1, p^*)} \leq \|\delta_\pi^* \phi\|_{(0, p^*)}, \quad (4.1)$$

holds for every $\phi \in \ker d_\pi$.

Proof. Let $\phi : S \rightarrow X^*$. Then

$$\begin{aligned} \kappa_{p^*}(S, X^*) \|\overline{D}\phi\|_{(2,p^*)} &= \kappa_{p^*}(S, X^*) 2^{\frac{1}{p^*}} \|\nabla\phi\|_{\ell_{p^*}(\mathcal{E}, X)} \\ &\geq 2^{\frac{1}{p^*}} \|\phi - \overline{M}\phi\|_{(1,p^*)} \\ &\geq 2^{\frac{1}{p^*}} \left(\|\phi\|_{(1,p^*)} - \|\overline{M}\phi\|_{(1,p^*)} \right) \end{aligned}$$

Since \overline{D} is an isometry on $\ker d_{\overline{\pi}}$,

$$2^{\frac{1}{p^*}} \|\phi\|_{(1,p^*)} - \kappa_{p^*}(S, X^*) \|\phi\|_{(1,p^*)} \leq 2^{\frac{1}{p^*}} \|\overline{M}\phi\|_{(1,p^*)},$$

which, by lemma 2.8, becomes

$$\left(1 - 2^{-\frac{1}{p^*}} \kappa_{p^*}(S, X^*) \right) \|\phi\|_{(1,p^*)} \leq \frac{1}{2} \|\delta_{\overline{\pi}}^* \phi\|_{(0,p^*)}.$$

□

Remark 4.2. The above inequality does not reduce to the one in [39] in the case $X = L_2$ and $p = 2$, even though in both cases the constant is non-zero if $\kappa_2(S, \mathbb{R}) < \sqrt{2}$. For $X = L_2$ and $p = 2$, Theorem 4.1 gives a strictly smaller lower estimate for the norm of the operator $\delta_{\overline{\pi}}^*$. Indeed, in that case the estimate obtained using spectral methods is

$$\sqrt{2(2 - \kappa_2(S, L_2)^2)} \|\phi\|_{(1,2)} \leq \|\delta_{\overline{\pi}}^* \phi\|_{(2,2)}.$$

This difference is a consequence of the fact that in the case $p = 2$ and $X = L_2$ we can apply the Pythagorean theorem instead of the triangle inequality in the first sequence of inequalities.

A similar inequality as in Theorem 4.1 holds for $\kappa_p(S, X)$ and the norm of $\delta_{\overline{\pi}}^*$. The above inequality can be now used to show that sufficiently small constants in Poincaré inequalities on the link graph imply fixed point properties.

4.2 Proof of the main theorem

Theorem 1.1. *Let X be a reflexive Banach space and let G be a group generated by a finite, symmetric set S , not containing the identity element. If the link graph $\mathcal{L}(S)$ is connected and for some $1 < p < \infty$ the associated Poincaré constants satisfy*


$$\max \left\{ 2^{-\frac{1}{p}} \kappa_p(S, X), 2^{-\frac{1}{p^*}} \kappa_{p^*}(S, X^*) \right\} < 1,$$

then

$$H^1(G, \pi) = 0,$$

for every isometric representation π of G on X .

Proof. We have the following dual diagrams:



For the purposes of this proof we will view, abusing the notation, δ_{π} as an operator $C^{(0,p)}(G, \pi) \rightarrow \ker d_{\pi}$. The natural injection of $\ker d_{\pi}$ into $C_{-}^{(1,p)}(G, \pi)$ will be denoted by i . Consequently, the formula (2.1) expresses the composition $\delta_{\pi}^* \circ i^*$ in this notation, as on the diagram. Similar notation is used on the dual level, with \bar{i} denoting the inclusion of $\ker d_{\bar{\pi}}$ into $C^{(1,p^*)}(G, \bar{\pi})$.

By Theorem 4.1, if $2^{-\frac{1}{p}} \kappa_p(S, X) < 1$ we conclude that $\delta_{\bar{\pi}}^* \circ \bar{i}^*$ is injective with closed image when restricted to $\ker d_{\pi}$. In fact, this means that the composition $\delta_{\bar{\pi}}^* \circ \bar{i}^* \circ i$ is injective with closed image. In particular, $\bar{i}^* \circ i$ is injective with closed image, and thus its dual, $i^* \circ \bar{i} : \ker d_{\bar{\pi}} \rightarrow (\ker d_{\pi})^*$, is surjective.

A similar argument applied to $2^{-\frac{1}{p^*}} \kappa_{p^*}(S, X^*) < 1$ yields $\delta_{\pi}^* \circ i^* \circ \bar{i}$ is also injective with closed image, which implies that δ_{π}^* is injective with closed image on the image of $i^* \circ \bar{i}$. Since the latter is surjective, δ_{π}^* is injective with closed image on $(\ker d_{\pi})^*$. This on the other hand implies that δ_{π} is onto, which proves the theorem by Proposition 2.10. \square

Remark 4.3. Note that under the assumptions of Theorem 1.1, $(\ker d_{\pi})^*$ and $\ker d_{\bar{\pi}}$ are isomorphic (a similar fact holds for $\ker d_{\pi}$ and $(\ker d_{\bar{\pi}})^*$). These spaces are closely related to the spaces of cocycles for the given representations.

It is an interesting question, for which isometric representations is $i^* \circ \bar{i}$ automatically an isomorphism, or at least is surjective. This property would eliminate, for such repre-

sentations, the need to use the inequality $2^{-\frac{1}{p}\kappa_p} < 1$, which is necessary in the above proof. A special case is discussed and applied in Section 7.2.

Remark 4.4. Note that it is not clear whether the above method can be extended to subspaces of $Y \subseteq X$. This would require estimating $\kappa_p(S, X)$ for some p , together with $\kappa_{p^*}(S, X^*)$, where Y^* is a quotient of X^* .

5 \tilde{A}_2 groups

In this section we apply Theorem 1.1 to specific groups and Banach spaces. In [7] the authors studied a family of groups $\{G_q\}$ called the A_2 -groups. These groups were introduced in [6], see also [4] for a detailed discussion. The group G_q has a presentation, whose associated link graph $\mathcal{L}(S)$ is the incidence graph of a finite projective plane $\mathbf{P}^2(\mathbb{F}_q)$ (here, q is a power of a prime number). Spectra of such graphs, with weight $\omega \equiv 1$, were computed by Feit and Higman [11], see also [4, 39]. It follows that

$$\kappa_2(S, \mathbb{R}) = \left(1 - \frac{\sqrt{q}}{q+1}\right)^{-\frac{1}{2}}.$$

In general, any estimates of p -Poincaré constants are difficult to obtain. In our case, the link graphs are finite graphs and we can use norm inequalities and a version of 3.0.5 to give the necessary estimates.

Theorem 5.1. *For each $q = k^n$ for some $n \in \mathbb{N}$ and prime number k we have*

$$H^1(G_q, \pi) = 0$$

for all

$$2 \leq p < \frac{\ln(q^2 + q + 1) + \ln(q + 1)}{\frac{1}{2} \ln(2(q^2 + q + 1)(q + 1)) - \ln(2) - \ln\left(\sqrt{1 - \frac{\sqrt{q}}{q+1}}\right)}$$

and for any isometric representation π of G_q on $L_p(Y, \mu)$ for any measure space.

Proof. We proceed by estimating $\kappa_p(S, L_p)$ and applying Theorem 1.1. Recall that given the space $\ell_p(\Omega)$ for $2 \leq p$, where the set Ω is finite, the following norm inequalities hold,

$$\|f\|_{\ell_p(\Omega)} \leq \|f\|_{\ell_2(\Omega)} \leq (\#\Omega)^{\frac{1}{2} - \frac{1}{p}} \|f\|_{\ell_p(\Omega)},$$

where $\|f\|_{\ell_p(\Omega)} = \left(\sum_{x \in \Omega} |f(x)|^p\right)^{\frac{1}{p}}$. Since the degree of the incidence graphs of finite projective planes is constant and equal to $q + 1$ we obtain for $f : S \rightarrow \mathbb{R}$ satisfying $Mf = 0$,

$$\begin{aligned} \|f\|_{(1,p)} &= (q+1)^{\frac{1}{p}} \|f\|_{\ell_p(S,X)} \\ &\leq (q+1)^{\frac{1}{p} - \frac{1}{2}} \|f\|_{(1,2)} \\ &\leq (q+1)^{\frac{1}{p} - \frac{1}{2}} \kappa_2(S, L_2) \|\nabla f\|_{\ell_2(\mathcal{E}, X)} \\ &\leq (q+1)^{\frac{1}{p} - \frac{1}{2}} \kappa_2(S, L_2) \left(\frac{\omega(E)}{2}\right)^{\frac{1}{2} - \frac{1}{p}} \|\nabla f\|_{\ell_p(\mathcal{E}, X)}. \end{aligned}$$

For each q we have $\omega(E) = 2(q^2 + q + 1)(q + 1)$, which gives the inequality

$$\begin{aligned} 2^{-\frac{1}{p}} \kappa_2(S, L_p) &\leq 2^{-\frac{1}{p}} (q + 1)^{\frac{1}{p} - \frac{1}{2}} \kappa_2(S, L_2) \left(\frac{\omega(E)}{2} \right)^{\frac{1}{2} - \frac{1}{p}} \\ &= 2^{-\frac{1}{p}} \left(\sqrt{1 - \frac{\sqrt{q}}{q + 1}} \right)^{-1} (q^2 + q + 1)^{\frac{1}{2} - \frac{1}{p}}. \end{aligned}$$

Bounding the above quantity by 1 from above gives

$$p < \frac{2 \ln(2(q^2 + q + 1))}{\ln(2(q^2 + q + 1)) - \ln \left(2 \left(1 - \frac{\sqrt{q}}{q + 1} \right) \right)}.$$

A similar norm estimate for $p^* \leq 2$, by virtue of the inequality

$$\|f\|_{\ell_2(\Omega)} \leq \|f\|_{\ell_{p^*}(\Omega)} \leq (\#\Omega)^{\frac{1}{p} - \frac{1}{2}} \|f\|_{\ell_2(\Omega)},$$

yields

$$p^* > \frac{2 \ln((q + 1)(q^2 + q + 1))}{\ln(2(q^2 + q + 1)(q + 1)) + \ln \left(2 \left(1 - \frac{\sqrt{q}}{q + 1} \right) \right)}$$

Simplifying and comparing p and $\frac{p^*}{p^* - 1}$ we obtain the claim. \square

Remark 5.2. The same argument gives a similar conclusion for the Banach space $X = (\bigoplus X_i)_p$, the ℓ_p -sum in which X_i is finite-dimensional with a norm sufficiently, and uniformly in i , close to the Euclidean norm. We leave the details to the reader.

Remark 5.3. The largest value of p in Theorem 5.1 is approximately 2.106 attained for $q = 13$. As q increases to infinity the values of p , for which cohomology vanishes, converge to 2 from above.

Remark 5.4. Although our estimate of the constant in the p -Poincaré inequality is not expected to be optimal, other interpolation methods do not seem to yield significantly better constants. For instance, Matousek’s interpolation method for p -Poincaré inequalities [22] gives a constant strictly greater than $2^{1/p}$ for any $p \geq 2$, since it emphasizes independence of dimension and is much better suited to deal with sequences of graphs (e.g. expanders).

Recall that the Banach-Mazur distance $d_{BM}(x, y)$ between two Banach spaces is the infimum of the set of numbers L , for which there exists an isomorphism $T : X \rightarrow Y$ satisfying $\|x\|_X \leq \|T_x\|_Y \leq L\|x\|_X$. Another consequence of Theorem 1.1 is that we obtain vanishing of cohomology for representations on Banach spaces, whose Banach-Mazur distance from the Hilbert space is controlled. We phrase this property in terms of uniformly bounded representations.

Theorem 5.5. *Let G_q be an \tilde{A}_2 -group and π be a uniformly bounded representation of G_q on a Hilbert space H , satisfying*

$$\sup_{g \in G} \|\pi_g\| < \sqrt{2 \left(1 - \frac{\sqrt{q}}{q+1}\right)}.$$

Then $H^1(G, \pi) = 0$.

Proof. Let $\|v\|' = \sup_{g \in G} \|\pi_g v\|$. Then $\|\cdot\|'$ is a norm and π is an isometric representation on $X = (H, \|\cdot\|')$. The identity is an isomorphism $\mathcal{I} : X \rightarrow H$ with $L = \sqrt{2 \left(1 - \frac{\sqrt{q}}{q+1}\right)}$, and $L\mathcal{I} : X^* \rightarrow H$ is an isomorphism with the same property. The estimate now follows by letting $p = 2$ and using the relation between $\kappa_2(S, X)$, $\kappa_2(S, X^*)$ and $\kappa_2(S, H)$, described in 3.0.5. \square

A similar fact (with appropriate constants) holds for L_p spaces, for the range of p as in the previous theorem.

6 Hyperbolic groups

In this section we discuss the consequences of 1.1 in the case of random hyperbolic groups. In [39] Żuk used spectral methods to show that many random groups have property (T) with overwhelming probability. A detailed account was recently provided in [20]. We sketch the strategy of the proof and generalize it to L_p -spaces.

In [13] it was shown that for a certain random graphs on n vertices of degree deg there exists a constant such that for any $\varepsilon > 1$ we have

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\kappa_2(S, \mathbb{R}) \leq \left(1 - \left(\frac{\sqrt{2 \text{deg}} (\text{deg} - 1)^{\frac{1}{4}}}{\text{deg}} + \frac{\varepsilon}{\text{deg}} \right) \right) \right) \rightarrow 1. \quad (6.1)$$

In [39] a modified link graph, denoted $L'(S)$, with multiple edges was considered. $L'(S)$ decomposes into random graphs as above and it is shown, using the above estimate, that it has a spectral gap strictly large than $1/2$ with probability 1. In our setting, the modified link graph $L'(S)$ can be viewed as a link graph with an admissible weight $\omega(s, t)$, which is defined to be the number of edges connecting s and t . Thus we can apply Theorem 1.1. Recall that in the Gromov model $\mathcal{G}(n, l, d)$ for random groups one chooses a density $0 < d < 1$ and considers a group given by a generating set S of cardinality n and $(2n-1)^{ld}$ relations of length l , chosen at random, letting l increase to infinity.

Theorem 6.1 ([39]; see also [20] for a detailed proof). *Let G be a random group in the density model, where $1/3 < d < 1/2$. Then, with probability 1, G is hyperbolic and there exists a group Γ and a homomorphism $\phi : \Gamma \rightarrow G$ with the following properties:*

1. Γ has a generating set S , whose link graph satisfies $2^{-1/2} \kappa_2(S, L_2) < 1$,
2. $\phi(\Gamma)$ is of finite index in G .

Given the above, we apply similar norm inequalities as in the case of \tilde{A}_2 -groups to the link graph of Γ and, as before, obtain fixed point properties for affine isometric actions of the group Γ on L_p for certain $p > 2$. For any given $p > 1$, the property of having vanishing cohomology $H^1(G, \pi)$ for *all* isometric representations π on L_p -spaces passes to quotients and from finite index subgroups to the ambient group. We thus have

Theorem 6.2. *With the assumptions of the previous theorem, with probability 1, Theorem 1.1 applies to hyperbolic groups. More precisely, let G, Γ and ϕ be as above, and let $\mathcal{L}(S) = (\mathcal{V}, \mathcal{E})$ denote the link graph of Γ . Then $H^1(G, \pi) = 0$ for every isometric representation π of G on L_p for*

$$p < \min \{p_0, \bar{p}_0^*\},$$

where

$$p_0 = \frac{\ln \deg_\omega - \ln(2\#\mathcal{E})}{\frac{1}{2} \ln \left(\frac{\deg_\omega}{\#\mathcal{E}} \right) - \ln \kappa_2(S, \mathbb{R})} \quad \text{and} \quad \bar{p}_0 = \frac{\ln(\#\mathcal{V} \deg_\omega) - \ln 2}{\frac{1}{2} \ln(\#\mathcal{V} \deg_\omega) - \ln \kappa_2(S, \mathbb{R})}.$$

We also have an estimate the norms of uniformly bounded representations to which cohomological vanishing can be extended for random hyperbolic groups.

Theorem 6.3. *Let G be a hyperbolic group in the Gromov model as above with $\frac{1}{3} < d < \frac{1}{2}$ and π be a uniformly bounded representation of G on a Hilbert space H , satisfying*

$$\sup_{g \in G} \|\pi_g\| < \frac{\sqrt{2}}{\kappa_2(S, \mathbb{R})},$$

Then $H^1(G, \pi) = 0$.

In other words, the $H^1(G, \pi)$ vanishes with probability 1 for representations bounded by $\sqrt{2}$.

We remark, that in (6.1), $\kappa_2(S, \mathbb{R})$ tends to 1 as $\deg \rightarrow \infty$. Thus the above upper bound on the norm of the representation is $\sqrt{2}$ with probability 1. On the other hand, Shalom showed that the $\text{Sp}(n, 1)$ has non-trivial cohomology with respect to some uniformly bounded representations (unpublished). The same property for hyperbolic groups is conjectured by Shalom and our Theorem 6.3 shows that one can extend property (T) to class of representations.

We also note that M. Cowling proposed to define a numerical invariant of a hyperbolic group by setting $\inf \{ \sup_{g \in G} \|\pi_g\| : H^1(G, \pi) \neq 0 \}$. Theorem 6.3 gives a uniform lower bound of $\sqrt{2}$ on such an invariant, with probability 1, for hyperbolic groups in the Gromov model.

Theorem 6.2 brings us to another interesting connection. Pansu [31] defined a quasi-isometry invariant of a hyperbolic group, called the conformal dimension, to be the number

$$\text{confdim}(\partial G) = \inf \{ \dim_{\text{Haus}}(\partial G, d) : d \text{ quasi-conformally equivalent to } d_v \},$$

where \dim_{Haus} denotes the Hausdorff dimension, ∂G denotes the Gromov boundary of the hyperbolic group G and d_v denotes any visual metric on ∂G . We refer to [18] for a brief overview of conformal dimension of boundaries of hyperbolic groups. Bourdon and Pajot [5] showed that a hyperbolic group acts by affine isometries without fixed points on $L_p(G)$ for p greater than the conformal dimension of ∂G . Combining this with vanishing of cohomology as studied here we see, that if $H^1(G, \pi)$ vanishes for all isometric representations on L_p then

$$p \leq \text{confdim}(\partial G).$$

Gromov [16, 9.B (g)] and Pansu [31, IV.b] posed the question of estimating the conformal dimension of random hyperbolic groups. Using Theorem 1.1 we obtain such estimates.

Corollary 6.4. *With the assumptions and notation of Theorem 6.2,*

$$\text{confdim}(\partial G) \geq \min \{p_0, \bar{p}_0^*\}.$$

Finally, as mentioned in the introduction, the above facts show that the method of Poincaré inequalities cannot in general give vanishing of cohomology as studied in this paper, for all $2 < p < \infty$. In addition, we have the following quantitative statement about Poincaré constants.

Corollary 6.5. *For any hyperbolic group G and any generating set S not containing the identity element, the Poincaré constants on the link graph associated to S satisfy*

$$\kappa_p(S, L_p) \geq 2^{\frac{1}{p}} \quad \text{or} \quad \kappa_{p^*}(S, L_{p^*}) \geq 2^{\frac{1}{p^*}},$$

for $p > \text{confdim}(\partial G)$.

7 Other Applications

7.1 Actions on the circle

Fixed point properties for the spaces L_p , $p > 2$, can be applied to studying actions on the circle, by applying the vanishing of cohomology to the L_p -Liouville cocycle. In [26] the following theorem was proved.

Theorem 7.1. *Let G be a discrete group, such that $H^1(G, \pi) = 0$ for every isometric representation of G on L_p for some $p > 2$. Then for every $\alpha > \frac{1}{p}$ every homomorphism $h : G \rightarrow \text{Diff}_+^{1+\alpha}(S^1)$ has finite image.*

Combining this result with, for instance, Theorem 5.1 we obtain

Corollary 7.2. *Let q be a power of a prime number and G_q be the corresponding \tilde{A}_2 group. Then every homomorphism $h : G \rightarrow \text{Diff}_+^{1+\alpha}(S^1)$ has finite image for*

$$\alpha > \frac{\frac{1}{2} \ln(2(q^2 + q + 1)(q + 1)) - \ln(2) - \ln \left(\sqrt{1 - \frac{\sqrt{q}}{q + 1}} \right)}{\ln(q^2 + q + 1) + \ln(q + 1)}.$$

7.2 The finite–dimensional case and the p -Laplacian

Let $1 < p < \infty$. The p -Laplacian Δ_p is an operator $\Delta_p : \ell_p(V) \rightarrow \ell_p(V)$, defined by the formula

$$\Delta_p f(x) = \sum_{x \sim y} (f(x) - f(y))^{[p]} \omega(x, y),$$

for $f : V \rightarrow \mathbb{R}$, where $a^{[p]} = |a|^{p-1} \text{sign}(a)$. The p -Laplacian reduces to the standard discrete Laplacian for $p = 2$, and is non-linear when $p \neq 2$. The p -Laplacian is of great importance in the study of partial differential equations. Its discrete version was studied e.g. in [1, 34]

A real number λ is an eigenvalue of the p -Laplacian Δ_p if there exists a function $f : V \rightarrow \mathbb{R}$ satisfying

$$\Delta_p f = \lambda f^{[p]}.$$

The eigenvalues of the p -Laplacian are difficult to compute in the case $p \neq 2$, due to non-linearity of Δ_p , see [15] for explicit estimates. Define

$$\lambda_1^{(p)}(\Gamma) = \inf \left\{ \frac{\sum_{x \in V} \sum_{y \sim x} |f(x) - f(y)|^p \omega(x, y)}{\inf_{\alpha \in \mathbb{R}} \sum_{x \in V} |f(x) - \alpha|^p \deg_\omega(x)} \right\}, \quad (7.1)$$

with the infimum taken over all $f : V \rightarrow \mathbb{R}$ such that f is not constant. $\lambda_1^{(p)}$ is the smallest positive eigenvalue of the discrete p -Laplacian Δ_p or the p -spectral gap.

We now apply an estimate similar to the one in Corollary 2.11, to finite quotients of groups. Let G be a finitely generated group and consider a homomorphism $h : G \rightarrow H$, where H is a finite group. Let $p > 1$ and let $\ell_p^0(H)$ denote the subspace of $\ell_p(H)$ consisting of those functions, which sum to 0.

We can identify the dual $\ell_p^0(H)^*$ with the space $\ell_{p^*}^0(H)$, with the norm

$$\|f\| = \inf_{\alpha \in \mathbb{R}} \|f - \alpha\|_{p^*}.$$

We will use our results to estimate the p^* -spectral gap for this norm on the Cayley graph of H .

Let $X = \ell_p^0(H)^*$ be equipped with the adjoint of the left regular representation λ on $\ell_p(H)$, restricted to $X^* = \ell_p^0(H)$. We have

$$\kappa_p(S, X^*) \leq \kappa_p(S, \ell_p(H)) = \kappa_p(S, \mathbb{R}).$$

Computing the Poincaré constant of the link graph for the norm of X is not straightforward. However, following the strategy outlined in Remark 4.3, we will show that we can bypass this condition. In order to do this we need to show that $i^* \circ \bar{i}$ is onto. In fact, a stronger statement is true.

Lemma 7.3. *Under the above assumptions, the map $i^* \circ \bar{i} : \ker d \rightarrow (\ker \bar{d})$, is an isomorphism.*

Proof. We can view X and X^* as having the same underlying vector space (real-valued functions $f : H \rightarrow X$ with mean value 0), equipped with two different norms. Similarly,

$C_-^{(1,p^*)}(G, \pi)$ and $C_-^{(1,p)}(G, \bar{\pi})$ also have the same underlying vector space, equipped with two different norms. The adjoint $\bar{\lambda}$ of the left regular representation, coincides with λ . For that reason $\ker d_\pi$ and $\ker d_{\bar{\pi}}$ describe the same vector subspace. The claim follows from the fact that all the spaces involved are finite-dimensional and complemented. \square

Now, since the representation of G on X does not have invariant vectors and δ_π is onto $\ker d_\pi$, we can conclude, by the Open Mapping Theorem, that δ_π in fact induces an isomorphism between $C^{(0,p^*)}(G, \pi)$ and $\ker d_\pi$. It follows from Theorem 4.1, that

$$2 \left(1 - 2^{-\frac{1}{p}} \kappa_p(S, \mathbb{R}) \right) \|f\|_{(0,p^*)} \leq \|\delta_\pi f\|_{1,p^*}.$$

Since $f \in \ell_p^0(H)$, this gives

$$\left(2 \left(1 - 2^{-\frac{1}{p}} \kappa_p(S, \mathbb{R}) \right) \right)^{p^*} \|f\|_X^{p^*} \leq \sum_{s \in S} \|f - \lambda_s f\|_X^{p^*} \frac{\deg_\omega(s)}{\omega(E)}.$$

Since $\|v\|_X \leq \|v\|_{\ell_p(H)}$, this yields

$$\begin{aligned} \left(2 \left(1 - 2^{-\frac{1}{p}} \kappa_p(S, \mathbb{R}) \right) \right)^{p^*} \inf_{c \in \mathbb{R}} \sum_{h \in H} |f(h) - c|^{p^*} \deg_\omega(h) \\ \leq \sum_{h \in H} \sum_{g \sim h} |f(h) - f(g)|^{p^*} \frac{\deg_\omega(g^{-1}h)}{\omega(E)}. \end{aligned}$$

(Note that $\deg_\omega(g^{-1}h)$ refer to $\mathcal{L}(S)$, not the Cayley graph of H .)

Corollary 7.4. *Let G be a group generated by a finite, symmetric set S not containing the identity element. If the link graph $\mathcal{L}(S)$ is connected and for some $1 < p < \infty$ the Poincaré constant satisfies*

$$2^{-\frac{1}{p}} \kappa_p(S, \mathbb{R}) < 1,$$

then

$$\lambda_1^{(p)} \geq 2 \left(1 - 2^{-\frac{1}{p}} \kappa_p(S, \mathbb{R}) \right)$$

on the Cayley graph of any finite quotient of G , for any weight $\omega(g, h) \geq \frac{\deg_\omega(g^{-1}h)}{\omega(E)}$.

Remark 7.5. A similar claim as in lemma 7.3 holds for any orthogonal representation which is also isometric on $\ell_p(H)$.

Acknowledgments.

I am grateful to Florent Baudier, Pierre-Emmanuel Caprace, Valerio Capraro, Tadeusz Januszkiwicz and Juhani Koivisto for helpful comments and to Masato Mimura and Andrzej Żuk for enjoyable conversations on related topics. I am also greatly indebted to Antoine Gournay and Alain Valette, who suggested several improvements and simplifications, in particular the use of the idempotent Q_π .

The author was partially supported by NSF grant DMS-0900874 and by the Foundation for Polish Science.

References

- [1] Amghibech, S.: Eigenvalues of the discrete p -Laplacian for graphs. *Ars Combin.* **67** (2003), 283–302
- [2] Bader, U., Furman, A., Gelfand, T., Monod, N.: Property (T) and rigidity for actions on Banach spaces. *Acta Math.* **198** (2007), no. 1, 57–105.
- [3] Ballmann, W., Świątkowski, J.: On L^2 -cohomology and property (T) for automorphism groups of polyhedral cell complexes. *Geom. Funct. Anal.* **7** (1997), no. 4, 615–645.
- [4] Bekka, B., de la Harpe, P., Valette, A.: Kazhdan’s property (T). *New Mathematical Monographs*, 11. Cambridge University Press, Cambridge, 2008.
- [5] Bourdon, M., Pajot, H.: Cohomologie l_p et espaces de Besov. *J. Reine Angew. Math.* **558** (2003), 85–108.
- [6] Cartwright, D. I., Mantero, A. M., Steger, T., Zappa, A.: Groups acting simply transitively on the vertices of a building of type \tilde{A}_2 . I. *Geom. Dedicata* **47** (1993), no. 2, 143–166.
- [7] Cartwright, D. I., Młotkowski, W., Steger, T.: Property (T) and \tilde{A}_2 groups. *Ann. Inst. Fourier (Grenoble)* **44** (1994), no. 1, 213–248.
- [8] Chatterji, I., Druţu, C., Haglund, F.: Kazhdan and Haagerup properties from the median viewpoint. *Adv. Math.* **225** (2010), no. 2, 882–921.
- [9] Dymara, J., Januszkiewicz, T.: Cohomology of buildings and their automorphism groups. *Invent. Math.* **150** (2002), no. 3, 579–627.
- [10] Ershov, M., Jaikin-Zapirain, A., Property (T) for noncommutative universal lattices. *Invent. Math.* **179** (2010), no. 2, 303–347.
- [11] Feit, W., Higman, G.: The nonexistence of certain generalized polygons. *J. Algebra* **1** 1964 114–131.
- [12] Fisher, D., Margulis, G.: Almost isometric actions, property (T), and local rigidity. *Invent. Math.* **162** (2005), no. 1, 1980.
- [13] Friedman, J.: Relative expanders or weakly relatively Ramanujan graphs. *Duke Math. J.* **118** (2003), no. 1, 19–35.
- [14] Garland, H.: p -adic curvature and the cohomology of discrete subgroups of p -adic groups. *Ann. of Math. (2)* **97** (1973), 375–423.
- [15] Grigorchuk, R. I., Nowak, P. W.: Diameters, distortion, and eigenvalues. *European J. Combin.* **33** (2012), no. 7, 1574–1587.

- [16] Gromov, M.: Asymptotic invariants of infinite groups. Geometric group theory, Vol. 2 (Sussex, 1991), 1295, London Math. Soc. Lecture Note Ser., 182, Cambridge Univ. Press, Cambridge, 1993.
- [17] Izeki, H., Nayatani, S.: Combinatorial harmonic maps and discrete-group actions on Hadamard spaces. *Geom. Dedicata* **114** (2005), 147–188.
- [18] Kapovich, I., Benakli, N.: Boundaries of hyperbolic groups. Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001), 3993, *Contemp. Math.*, 296, Amer. Math. Soc., Providence, RI, 2002.
- [19] Kassabov, M.: Subspace arrangements and property T. *Groups Geom. Dyn.* **5** (2011), no. 2, 445–477.
- [20] Kotowski, M., Kotowski, M.: Random groups and property (T): Żuk’s theorem revisited. *J. Lond. Math. Soc. (2)* **88** (2013), no. 2, 396–416.
- [21] Lafforgue, V.: Un renforcement de la propriété (T). *Duke Math. J.* **143** (2008), no. 3, 559–602.
- [22] Matoušek, J.: On embedding expanders into l_p spaces. *Israel J. Math.* **102** (1997), 189–197.
- [23] Mimura, M.: Fixed point properties and second bounded cohomology of universal lattices on Banach spaces. *J. Reine Angew. Math.* **653** (2011), 115–134.
- [24] Naor, A., Silberman, L.: Poincaré inequalities, embeddings, and wild groups. *Compos. Math.* **147** (2011), no. 5, 1546–1572.
- [25] Navas, A.: Actions de groupes de Kazhdan sur le cercle. *Ann. Sci. École Norm. Sup. (4)* **35** (2002), no. 5, 749–758.
- [26] Navas, A.: Reduction of cocycles and groups of diffeomorphisms of the circle. *Bull. Belg. Math. Soc. Simon Stevin* **13** (2006), no. 2, 193205.
- [27] Nica, B.: Proper isometric actions of hyperbolic groups on L^p -spaces. *Compos. Math.* **149** (2013), no. 5, 773–792.
- [28] Nowak, P. W.: Group actions on Banach spaces and a geometric characterization of a-T-menability. *Topology Appl.* **153** (2006), no. 18, 3409–3412.
- [29] Nowak, P. W.: Group actions on Banach spaces and a geometric characterization of a-T-menability. (corrected version). [arXiv:math/0404402v3](https://arxiv.org/abs/math/0404402v3)
- [30] Nowak, P. W.: Group Actions on Banach Spaces. *Handbook of Group Actions*, ed.: by L. Ji, A. Papadopoulos, S.-T. Yau, to appear.
- [31] Pansu, P.: Dimension conforme et sphère à l’infini des variétés à courbure négative. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **14** (1989), no. 2, 177–212.

- [32] Pansu, P.: Cohomologie L_p : invariance sous quasiisométrie. Preprint, 1995.
- [33] Pansu, P.: Formules de Matsushima, de Garland et propriété (T) pour des groupes agissant sur des espaces symétriques ou des immeubles Bull. Soc. Math. France 126 (1998), no. 1, 107–139.
- [34] Takeuchi, H.: The spectrum of the p -Laplacian and p -harmonic morphisms on graphs. Illinois J. Math. **47** (2003), no. 3, 939–955.
- [35] Wang, M.-T.: A fixed point theorem of discrete group actions on Riemannian manifolds. J. Differential Geom. **50** (1998), no. 2, 249–267.
- [36] Wang, M.-T.: Generalized harmonic maps and representations of discrete groups. Comm. Anal. Geom. **8** (2000), no. 3, 545–563.
- [37] Yu, G.: Hyperbolic groups admit proper affine isometric actions on l^p -spaces. Geom. Funct. Anal. **15** (2005), no. 5, 1144–1151.
- [38] Żuk, A.: La propriété (T) de Kazhdan pour les groupes agissant sur les polyèdres. C. R. Acad. Sci. Paris Sr. I Math. **323** (1996), no. 5, 453–458.
- [39] Żuk, A.: Property (T) and Kazhdan constants for discrete groups. Geom. Funct. Anal. **13** (2003), no. 3, 643–670.